

Letter to the Editor

Phenotypic Discordance in a Family With Monozygotic Twins and Non-Syndromic Cleft Lip and Palate

To the Editor:

Despite considerable research, the cause of non-syndromic cleft lip with or without cleft palate (NSCLP) is still an enigma. Case-control and cohort studies have searched for environmental factors that might influence the development of this common malformation, such as maternal cigarette smoking [Saxén, 1974; Evans et al., 1979; Ericson et al., 1979; Shiono et al., 1986; Khoury et al., 1987, 1989; Werler et al., 1990; Hwang et al., 1995], periconceptional supplementation of folic acid and multivitamins [Tolarová and Harris, 1995; Shaw et al., 1995], agricultural chemical use [Gordon and Shy, 1981; Nurminen et al., 1995], and place of residence [Christensen et al., 1995], among others. However, these studies are subject to numerous biases [Khoury et al., 1992; Hemminki et al., 1995; Nurminen, 1995; Lie, 1995], and their results have often been contradictory and inconclusive.

There is a well-known genetic component to the cause of NSCLP, but its specific nature remains uncertain [Juriloff, 1993]. Researchers performing complex segregation analyses have proposed a single gene of major effect for NSCLP, with autosomal recessive [Chung et al., 1986; Marazita et al., 1992], or dominant [DePaepe, 1989; Temple et al., 1989; Hecht et al., 1990; Ray et al., 1993] inheritance, but always with incomplete penetrance. Several groups are currently engaged in linkage studies intended to identify major gene[s] for NSCLP. Various candidate regions have been reported to show linkage or allelic association with NSCLP: F13A on chromosome 6 [Eiberg et al., 1987], transforming growth factor- α [TGF α] on chromosome 2 [Ardinger et al., 1989], retinoic acid receptor- α [RAR α] on chromosome 17 [Chenevix-Trench et al., 1992], MFD38 on chromosome 4 [Beiraghi et al., 1994], chromosomal region 6p24 [Davies et al., 1995], and BCL3 on chromosome 19 [Stein et al., 1995]. The lack of support by other studies [Hecht et al., 1991, 1993; Vintiner et al., 1992] has been addressed in detail by Farrall et al. [1993] and Murray [1995].

Received for publication December 14, 1995; revision received April 9, 1996.

Address reprint requests to Dr. Diego F. Wyszynski, Department of Epidemiology, School of Hygiene and Public Health, The Johns Hopkins University, 615 North Wolfe Street, Baltimore, MD 21205.

CLINICAL REPORT

Here we present a clinical case to generate interest in studies of recurrence risk for discordant MZ twins for CL/P, an area still unexplored in the literature. In 1994, as part of a linkage study, we met a NSCLP family with two affected members (Fig. 1). The proband, individual 8, was diagnosed by a medical geneticist as having a right cleft lip (CL) and a complete cleft palate (CP). The uvula was intact. The child was the product of an uncomplicated pregnancy, and his mother reported no exposure to teratogens during the first trimester of gestation. Multivitamin supplements were taken between the second and fifth months of pregnancy. Maternal and paternal age at time of delivery were 27 and 26 years, respectively. Birth and development were uncomplicated. The patient's mother and aunt are identical twins. His mother, individual 4 in Figure 1, was diagnosed as having a left CL with a complete CP. No uvula was present. Her sister, individual 5, was born with no oral cleft and an intact uvula. Review of the obstetric charts from this twin pregnancy showed no complications. Individual 7, brother of the twin sisters, had an isolated syndactyly between the third and fourth finger in one hand, enough to prevent wearing a ring, but this individual was otherwise normal.

In addition to the high degree of physical resemblance between the twins, monozygosity was confirmed by polymerase chain reaction (PCR) amplification and analysis of six polymorphic markers on different chromosomes [Eufinger et al., 1995] (data available upon request). The patient's mother and aunt had perfect agreement for all six markers.

Cases of MZ twins discordant for genetically determined disorders have been reported, e.g., Ullrich-Turner syndrome or Duchenne muscular dystrophy [Eufinger et al., 1993]. In these cases postzygotic mitotic nondisjunction or anaphase lags, as well as uneven X-chromosome inactivation are possible explanations [Gonsoulin et al., 1990; Perlman et al., 1990; Lupski et al., 1991]. Previous twin studies of NSCLP have reported concordance rates of 36% (range 33–78) for MZ twins and 1% [range 0–14.3] for DZ twins [Eufinger et al., 1993]. As MZ twins possess identical genomes, discordance or differing grades of clefting may be explained by exposure to potential teratogenic agents (i.e., respiratory viruses [Zhang and Cai, 1993]), a postzygotic mutation, or the effect of chance. This family is particularly interesting, since the affected MZ twin gave birth to an affected child, thus making in




Fig. 1. Pedigree of family 40. Individual 4 has a left cleft lip (CL) and a complete cleft palate (CP). Individual 8, the proband, has a right CL and a complete CP. Individual 7 has a webbed finger in a hand but he is otherwise normal.

utero environmental factors unlikely to be solely responsible for the genesis of these oral clefts. This raises the question of how best to counsel the unaffected sister (individual 5 in Fig. 1) as to her risk of having a child with a NSCLP: "Is it the same as her identical twin's risk (approximately 8–10 times higher than the risk of the general population) or lower (3–5 times), as she herself is 'unaffected'?" The risk for the unaffected sister to have a child with CL/P was given as 3–10% (potentially as high as that for her affected sister, though likely to be less). This estimate has been debated by other geneticists. While this family is being followed, additional twin studies, including families like the one presented here, may provide more appropriate estimates of risk.

ACKNOWLEDGMENTS

We are indebted to family 044 for participating in our study. We would like to acknowledge Mrs. Jennifer Simpson, RNC, for her help contacting the family and collecting the samples. Dr. David L. Duffy provided important comments on the final manuscript. Finally, we would like to thank Mrs. E. Anne Smith and Dr. Iain McIntosh for genotyping the samples. This study was supported in part by grant D01-DE10293 from the National Institutes of Health.

REFERENCES

Ardinger HH, Buetow KH, Bell GI, Bardach J, Vandemark DR, Murray JC (1989): Association of genetic variation of the transforming growth factor- α gene with cleft lip and palate. *Am J Hum Genet* 45:348–353.

Beiraghi S, Foroud T, Diouhy S, Bixler D, Conneally PM, Delozier-Blanchet D, Hodes M (1994): Possible localization of a major gene for cleft lip and palate to 4q. *Clin Genet* 46:255–256.

Chenevix-Trench G, Jones K, Green AC, Duffy DL, Martin NG (1992): Cleft lip with or without cleft palate: Associations with transforming growth factor alpha and retinoic acid receptor loci. *Am J Hum Genet* 51:1377–1385.

Christensen K, Schmidt MM, Væth M, Olsen J (1995): Absence of an environmental effect on the recurrence of facial-cleft defects. *New Engl J Med* 333:161–164.

Chung CS, Bixler D, Watanabe T, Koguchi H, Fogh-Andersen P (1986): Segregation analysis of cleft lip with or without cleft palate: A comparison of Danish and Japanese data. *Am J Hum Genet* 39: 603–611.

Davies AF, Stephens RJ, Olavesen MG, Heather L, Dixon MJ, Magee A, Flint F, Ragoussis J (1995): Evidence of a locus for orofacial clefting on human chromosome 6p24 and STS content map of the region. *Hum Molec Genet* 4:121–128.

DePaepe A (1989): Dominantly inherited cleft lip and palate. *Letter. J Med Genet* 26:794.

Eiberg H, Bixler D, Nielsen LS, Conneally PM, Mohr J (1987): Suggestion of linkage of a major locus for nonsyndromic orofacial cleft with F13A and tentative assignment to chromosome 6. *Clin Genet* 32:129–132.

Ericson A, Källén B, Westerholm P (1979): Cigarette smoking as an etiologic factor in cleft lip and palate. *Am J Obstet Gynecol* 135: 348–351.

Eufinger H, Rand S, Scholz W, Machtens E (1993): Clefts of the lip and palate in twins: Use of DNA fingerprinting for zygosity determination. *Cleft Palate-Craniofac J* 30:564–568.

Eufinger H, Rand SP, Schüte U (1995): Use of single- and multi-locus and polymerase chain reaction systems for zygosity determination: Clinical application in twins with clefts of the lip and palate. *Acta Genet Med Gemellol* 44:25–30.

Evans DR, Newcombe RG, Campbell H (1979): Maternal smoking habits and congenital malformations: A population study. *Br Med J* 2:171–173.

Farrall M, Buetow KH, Murray JC (1993): Resolving an apparent paradox concerning the role of TGFA in CL/P. *Letter. Am J Hum Genet* 52:434–436.

Gonsoulin W, Copeland KL, Carpenter RJ, Hughes MR, Elder FF (1993): Fetal blood sampling demonstrating chimerism in monozygotic twins discordant for sex and tissue karyotype (46,XY and 45,X). *Prenat Diagn* 10:25–28.

Gordon JE, Shy CM (1981): Agricultural chemical use and congenital cleft lip and/or palate. *Arch Environm Health* 36:213–220.

Hecht JT (1990): Dominantly inherited cleft lip and palate (Letter to the Editor). *J Med Genet* 27:597–598.

Hecht JT, Wang Y, Blanton SH, Michels VV, Daiger SP (1991): Cleft lip and palate: no evidence of linkage to transforming growth factor alpha. *Am J Hum Genet* 49:682–686.

Hecht JT, Wang Y, Connor B, Blanton SH, Daiger SP (1993): Nonsyndromic cleft lip and palate: No evidence of linkage to HLA or factor 13A. *Am J Hum Genet* 52:1230–1233.

Hemminki K, Lindbohm M-L, Kyrrönen P (1995): Validity aspects of exposure and outcome data in reproductive studies. *J Occup Environm Med* 37:903–907.

Hwang S-J, Beaty TH, Panny SR, Street NA, Joseph JM, Gordon S, McIntosh I, Francomano CA (1995): Association study of transforming growth factor alpha (TGF α) TaqI polymorphism and oral clefts: Indication of gene-environment interaction in a population-based sample of infants with birth defects. *Am J Epidemiol* 141: 629–636.

Juriloff DM (1993): Current status of genetic linkage studies of a major gene that causes CL(P) in mice: exclusion map. *J Craniofac Genet Dev Biol* 13:223–229.

Khoury MJ, Weinstein A, Panny S, Holtzman NA, Lindsay PK, Farrel K, Eisenberg M (1987): Maternal cigarette smoking and oral clefts: A population-based study. *Am J Pub Health* 77:623–625.

Khoury MJ, Gomez-Farias M, Mulinare J (1989): Does maternal cigarette smoking during pregnancy cause cleft lip and palate in offspring? *Am J Dis Children* 143:333–337.

Khoury MJ, James LM, Flanders WD, Erickson JD (1992): Interpretation of recurring weak associations obtained from epidemiologic studies of suspected human teratogens. *Teratology* 46:69–77.

Lie RT (1995): Flawed epidemiology of birth defects? *Letter. Lancet* 346:1037.

Lupski JR, Garcia CA, Zoghbi HY, Hoffman EP, Fenwick RG (1991): Discordance of muscular dystrophy in monozygotic female twins: Evidence supporting asymmetric splitting of the inner cell mass in

- a manifesting carrier of Duchenne dystrophy. *Am J Med Genet* 40: 354–364.
- Marazita ML, Hu D-N, Spence MA, Liu Y-E, Melnick M (1992): Cleft lip with or without cleft palate in Shanghai, China: Evidence for an autosomal major locus. *Am J Hum Genet* 51:648–653.
- Murray JC (1995): Face facts: Genes, environment, and clefts. *Am J Hum Genet* 57:227–232.
- Nurminen T (1995): Maternal pesticide exposure and pregnancy outcome. *J Occup Environm Med* 37:935–940.
- Perlman EJ, Stetten G, Tuck-Muller CM, Farber RA, Neuman WL, Blakemore KJ, Hutchins GM (1990): Sexual discordance in monozygotic twins. *Am J Med Genet* 37:551–557.
- Ray AK, Field LL, Marazita ML (1993): Nonsyndromic cleft lip with or without cleft palate in West Bengal, India: Evidence for an autosomal major locus. *Am J Hum Genet* 52: 1006–1011.
- Saxén I (1974): Cleft lip and palate in Finland: Parental histories, course of pregnancy and selected environmental factors. *Int J Epid* 3:263–270.
- Shaw GM, Lammer EJ, Wasserman CR, O'Malley CD, Tolarová MM (1995): Risks of orofacial clefts in children born to women using multivitamins containing folic acid periconceptionally. *Lancet* 346: 393–396.
- Shiono PH, Klebanoff MA, Berendes HW (1986): Congenital malformations and maternal smoking during pregnancy. *Teratology* 34: 65–71.
- Stein J, Mulliken JB, Stal S, Gasser DL, Malcolm S, Winter R, Blanton SH, Amos C, Seemanová E, Hecht J (1995): Nonsyndromic cleft lip with or without cleft palate: evidence of linkage to BCL3 in 17 multigenerational families. *Am J Hum Genet* 57:257–272.
- Temple K, Calvert M, Plint D, Thompson E, Pembrey M (1989): Dominantly inherited cleft lip and palate in two families. *J Med Genet* 26:386–389.
- Tolarová M, Harris J (1995): Reduced recurrence of orofacial clefts after periconceptional supplementation with high-dose folic acid and multivitamins. *Teratology* 51:71–78.
- Vintiner GM, Holder SE, Winter RM, Malcolm S (1992): No evidence of linkage between the transforming growth factor-alpha gene in families with apparently autosomal dominant inheritance of cleft lip and palate. *J Med Genet* 29:393–397.
- Werler MM, Lammer EJ, Rosenberg L, Mitchell A (1990): Maternal cigarette smoking during pregnancy in relation to oral clefts. *Am J Epidemiol* 132:926–932.
- Zhang J, Cai W-W (1993): Association of the common cold in the first trimester of pregnancy with birth defects. *Pediatrics* 92:559–563.

Diego F. Wyszynski

Department of Epidemiology
School of Hygiene and Public Health
The Johns Hopkins University
Baltimore, Maryland
Medical Genetics Branch
National Center for Human Genome Research
Bethesda, Maryland

Amy F. Lewanda

Department of Pediatrics
The Johns Hopkins Hospital
Baltimore, Maryland
Department of Medical Genetics
Children's National Medical Center
Washington, D.C.

Terri H. Beaty

Department of Epidemiology
School of Hygiene and Public Health
The Johns Hopkins University
Baltimore, Maryland